## (12)

## EUROPEAN PATENT APPLICATION

(1) Application number: 89302442.2

(a) Int. Cl.4: C12P 17/18 , //(C12P17/18, C12R1:465),A61K31:00

@ Date of filing: 13.03.89

- Priority: 14.03.88 GB 8805978
- Date of publication of application:
  20.09.89 Bulletin 89/38
- Designated Contracting States:
  AT BE CH DE ES FR GB GR IT LI LU NL SE
- Applicant: AMERICAN CYANAMID COMPANY
   One Cyanamid Plaza
   Wavne New Jersey 07470(US)
- Inventor: Rudd, Brian Arthur Michael 72 Exeter Road Rayners Lane Harrow Middlesex(GB) Inventor: Ramsay, Michael Vincent John 157 Kings Road South Harrow Middlesex(GB)
- Representative: Skalles, Humphrey John et al Frank B. Dehn & Co. Imperial House 15-19 Kingsway London WC2B 6UZ(GB)
- Process for the preparation of macrolide compounds.
- (F) S541 (LL-F28249) compounds of formula (I)

[wherein R¹ is a methyl, ethyl or isopropyl group; R² is a hydrogen atom or a group  $OR^5$  (where  $OR^5$  is a hydroxyl group or a substituted hydroxyl group) and  $R^3$  is a hydrogen atom, or  $R^2$  and  $R^3$  to gether with the carbon atom to which they are attached represent  $> 0 < 0 < C < Ch_2$  or > 0 < C > CR6 (where R¹ represents a hydrogen atom,  $C_{1-2}$  alkyl group or a  $C_{3-2}$  alkenyl group and the group  $> C < ROR^6$  is in the E configuration); and R¹ is a hydrogen atom or a methyl group) which comprises incubating a corresponding a suitable medium in the presence of a microorganism or an enzyme derived therefrom

o 0 333 404 A2

or a preparation derived from a microorganism containing the enzyme of interest capable of effecting the

#### PROCESS FOR THE PREPARATION OF MACROLIDE COMPOUNDS

This invention relates to a process for the production of macrolide compounds.

15

20

25

United Kingdom Patent Specification 2166436 and European Patent Specification 170006 describe a class of substances, which we have designated Antibiotics 5541, which may be prepared by fermentation of Antibiotics 5541 producing strains belonging to the species Streptomyces thermoarchaensis and Streptomyces cyaneogriseus noncyanogenus. Antibiotics 5541 have antibiotic and, in particular, anti-endoparasitic, anti-ectoparasitic, anti-tungal, insecticidal, nematicidal and acaricidal activity and are of special interest for use in agriculture, horticulture, and animal and human health.

We have now found a new process for the production of Antibiotics S541 compounds and chemical derivatives thereof.

Thus, according to one aspect of the invention we provide a process for the production of compounds of formula (I)

[wherein R¹ is a methyl, ethyl or isopropyl group; R² is a hydrogen atom or a group OR⁵ where OR⁵ is a hydroxyl group or a substituted hydroxyl group having up to 25 carbon atoms) and R¹ is a  $P_1$  for  $P_2$  for  $P_3$  for  $P_$ 

(wherein R¹, R² and R³ are as defined previously) in a suitable medium in the presence of a microorganism or an enzyme derived therefrom or a preparation derived from a microorganism containing the enzyme of interest capable of effecting the conversion. Suitable microorganisms and extracts thereof for use in the process according the invention may be identified by preliminary small scale tests designed to demonstrate ability of a microorganism or an extract thereof to convert compounds of formula (i) to compounds of formula (i). The formulation of the compounds of formula (i) may be confirmed by suitable chromatographic analysis (eg high performance iliquid s chromatography) of the reaction mixture.

We have found microorganisms of the genus <u>Streptomyces</u> and extracts thereof to be particularly suitable for use in the process according to the present invention.

Particular Streptomyces microorganisms for use in the process according to the invention include strains of Streptomyces thermoarchaensis, Streptomyces cyaneogriseus noncyanogenus, Streptomyces avermitilis and Streptomyces hygroscopicus subsp, aureolacrimosus. Particular examples of suitable strains include Streptomyces thermoarchaensis NCIB 12212 and 12213 (deposited 6th March 1988 in the permanent culture collection of the National Collections of Industrial and Marine Bacteria, Torry Research Station, Aberdeen, United Kingdom] and mutants of these strains, Streptomyces avermitilis ATCC 31272 and 31780 and Streptomyces hygroscopicus subsp, aureolacrimosus FERM P 1438.

Mutants of the above strains may arise spontaneously or may be produced by a variety of methods including those described in UK Patent Specification 2166436 and European Patent Specification 170006.

Suitable enzymes for use in the process according to the present invention may be derived from an extremely wide range of sources. The aforementioned Streptomyces microorganisms, however, represent a particularly suitable source of enzymes capable of converting compounds of formula (II) into compounds of formula (II).

In one embodiment of the process according to the invention, the conversion of a compound of formula (II) into a corresponding 5-OR\* compound of formula (I) may be effected by feeding the compound of formula (II) in a suitable solvent into a fermentation medium comprising the aforementioned microorganism in the presence of assimilable sources of carbon, nitroen and mineral salts.

Assimilable sources of carbon, nitrogen and minerals may be provided by either simple or complex nutrients. Sources of carbon will generally include glucose, maltose, starch, glycerol, molasses, dextrin, lactose, sucrose, fructose, carboxylic acids, amino acids, glycerides, alchols, alkanes and vegetable oils. Sources of carbon will generally comprise from 0.5 to 10% by weight of the fermentation medium.

Sources of nitrogen will generally include soya bean meal, com steep liquors, distillers solubles, yeast 30 extracts, cottonseed meal, peptones, ground nut meal, malt extract, molasses, casein, amino acid mixtures, ammonia (gas or solution), ammonium salts or nitrates. Urea and other amides may also be used. Sources of nitrogen will generally comprise from 0.1 to 10% by weight of the fermentation medium.

Nutrient mineral salts which may be incorporated into the culture medium include the generally used salts capable of yielding sodium, potassium, ammonium, iron, magnesium, zinc, nickel, cobalt manganese, su vanadium, chromium, calcium, copper, molybdenum, boron, phosphate, sulphate, chloride and carbonate

An antifoam may be present to control excessive foaming and added at intervals as required.

The compound of formula (II) in a solvent such as a water miscible organic solvent (eg an alcohol such as methanol or propan-2-ol, a diol such as propan-1,2-ol or butan-1,3-ol, a ketone such as acetone, a nitrile such as acetontifile, an ether such as tetrahydrofuran or dioxan, a substituted amide such as dimethylformamide or a dialkylsulphoxide such as dimethylsulphoxide) may be added at the beginning of the cultivation, or more usually, when the growth of the microorganism is under way, e.g. 2 to 4 days after the start of the cultivation.

Cultivation of the organism will generally be effected at a temperature of from 20 to 50°C, preferably from 25 to 40°C, and will desirably take place with aeration and agitation e.g. by shaking or stirring. The medium may initially be inoculated with a small quantity of a suspension of the sporulated microorganism but in order to avoid a growth lag a vegetative inoculum of the organism may be prepared by inoculating a small quantity of the culture medium with the spore form of the organism, and the vegetative inoculum obtained may be transferred to the fermentation medium, or, more preferably to one or more seed stages where further growth takes place before transfer to the principal fermentation medium. The fermentation will cenerally be carried out in the oH range 5.5 to 8.5, preferably 5.5 to 7.5.

Once the compound of formula (II) has been added to the culture, usually with gentle mixing, the cultivation is continued such that the desired product is accumulated. The presence of the product in the fermentation broth may be determined by monitoring extracts of the broth by high performance liquid chromatography, and uv spectroscopy at 238nm.

The product(s) may be isolated from the whole fermentation broth by conventional isolation and separation techniques as described in UK Patent Specifications 2166436 and 2176162.

In a further embodiment of the process according to the invention, the conversion of a compound of

formula (II) into a corresponding 5-OR\* compound of formula (I) may be effected by combining and incubating a compound of formula (II) in a suitable solvent (eg a water miscible organic solvent as previously defined) with a preparation of the enzyme of the invention, desirably in a buffer solution, at, for example, 0° to 60°, preferably 20° to 40° eg about 28°C. The reaction will generally be carried out in the 5 pH range 3.5 to 8.5 eq 5.5 to 7.5. If desired the reaction may be carried out in the presence of a cofactor such as NADH or NADPH. When the reaction is complete, le when the compound of formula (II) is no longer converted to the compound of the invention (as determined by monitoring extracts of the reaction mixture by high performance liquid chromatography and uv spectroscopy at 238nm) the product is recovered by conventional isolation and separation techniques as described in UK Patent Specifications 10 2166436 and 2176182.

The enzyme for use in the process of the present invention may be prepared, for example, by culture of a micrographism which produces the enzyme in a nutrient medium. Suitable nutrient media and fermentation conditions for the preparation of the enzyme include those previously described for the preparation of a compound of formula (I) from a compound of formula (II) In the presence of a 15 microorganism. The time at which the required enzymic activity reaches a maximum will, of course, vary according to the microorganism used and, hence, the optimum cultivation time is desirably determined independently for each strain employed.

For microproanisms where the enzyme is extracellular, the liquid culture medium or the filtrate after removal of whole cells may be used as a source of enzyme. Where the enzyme is cell-bound it may be 20 released for use by conventional methods such as sonication, grinding with glass beads, homogenisation, treatment with lytic enzymes or with detergents, after suspension of the cells in a suitable buffer,

The resulting preparation, either with or without removal of cell debris, may be used as a source of enzyme. It is preferred, however, to purify the enzyme further by conventional means. Batch or column chromatography with ion-exchange celluloses or affinity adsorbents or other adsorbents eq hydroxylapatite 25 may conveniently be employed. In addition, the enzyme may be concentrated or further purified by molecular sieve techniques eg ultrafiltration or salting out. In general, during purification procedures, it is desirable to maintain the pH within the range 3 to 11.

The enzyme may be employed in an immobilized form, eg by insolubilisation or entrappment thereof on or in a suitable matrix. Thus an extract of the enzyme may be bound or linked to an otherwise inert 30 inorganic or organic polymer, entrapped on or in a fibre, or on or in a membrane or polymer such as polyacrylamide gel, adsorbed on an ion-exchange resin, crosslinked with a reagent such as glutaraldehyde, or occluded in an envelope such as a bead. Immobilized enzymes may advantageously be employed both in batch processes, after which the enzyme may be reused, and continuous flow processes wherein substrates pass through a column containing the immobilized enzyme.

The group R5 when present in compounds of formula (I) may represent an acyl group e.g. a group of the formula B<sup>2</sup>CO- or B<sup>2</sup>OCO- or B<sup>2</sup>OCO- (where B<sup>2</sup> is an aliphatic, araliphatic or aromatic group, for example an alkyl, alkenyl, alkynyl, cycloalkyl, aralkyl or aryl group), a formyl group, a group R8 which is as defined above for R7, a group R9SO₂- (where R9 is a C1-4 alkyl or C6-10 aryl group), a silyl group, a cyclic or acyclic acetal group, a group -CO(CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R<sup>10</sup> (where R<sup>10</sup> is a hydrogen atom or a group as defined 40 above for R7 and n represents zero, 1 or 2) or a group R11R12NCO- (where R11 and R12 may each Independently represent a hydrogen atom or a C1-4 alkyl group).

35

Where R7 or R8 are alkyl groups, they may be for example C1-8 alkyl groups, e.g. methyl, ethyl, npropyl, i-propyl, n-butyl, i-butyl, t-butyl or n-heptyl which alkyl groups may also be substituted. Where R7 is a substituted alkyl group it may be substituted by, for example, one or more, eg two or three, halogen atoms (e.g. chlorine or bromine atoms), or a carboxy, C1-4 alkoxy (e.g. methoxy, ethoxy), phenoxy or silvloxy group. Where R8 is a substituted alkyl group it may be substituted by a cycloalkyl e.g. cyclopropyl

Where R7 and R8 are alkenyl or alkynyl groups, they preferably have 2-8 carbon atoms and where R7 and R8 are cycloalkyl groups, they may be for example C3-12 cycloalkyl, such as C3-7 cycloalkyl, eg cyclopentyl groups.

Where R7 and R8 are aralkyl groups, they preferably have 1-6 carbon atoms in the alkyl moiety, and the aryl group(s) may be carbocylic or heterocyclic and preferably contain 4-15 carbon atoms e.g. phenyl. Examples of such groups include phen C1-5 alkyl eg benzyl groups.

Where R7 and R8 are anyl groups, they may be carbocyclic or heterocyclic and preferably have 4-15 55 carbon atoms e.g. phenyl.

When R5 is a group R9SO2-, it may be for example a methylsulphonyl or p-toluenesulphonyl group.

Where R5 represents a cyclic acetal group, it may for example have 5-7 ring members as in the tetrahydropyranyl group.

When R<sup>5</sup> represents a sliyl group or R<sup>7</sup> contains a silyloxy substituent, the silyl group may carry three groups which may be the same or different, selected from alkyl, alkenyl, alkoxy, cycloalkyl, aralkyl, aryl and aryloxy groups. Such groups may be as defined above and particularly include methyl, t-butyl and phenyl groups. Particular examples of such sliyl groups are trimethylsilyl and t-butyldimethylsilyl.

When  $R^5$  represents a group -CO(CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>R<sup>10</sup>, it may for example be a group -COCO<sub>2</sub>R<sup>10</sup> or -COCH<sub>2</sub>CH<sub>2</sub>CO<sub>2</sub>R<sup>10</sup> where R<sup>10</sup> represents a hydrogen atom or a C<sub>1-4</sub> alkyl group (eg methyl or ethyl).

When R5 represents a group R'I'R'2NCO-, R'1 and R12 for example may each independently be a hydrogen atom or a methyl or ethyl group.

When R<sup>6</sup> represents a C<sub>1-3</sub> alkyl group it may be for example a methyl, ethyl, n-propyl, i-propyl, n-

When R6 regresents a C3-8 alkenyl group it may be for example an allyl group.

Preferred compounds of formula (I) which may be prepared according to the process of the present invention include compounds in which R' is isopropyl.

Important compounds of formula (I) which may be prepared according to the process of the present invention include those in which R² represents a hydroxy or ethoxy group and R³ represents a hydrogen atom; or R² and R³ each represents a hydrogen atom; or R² and R³ together with the carbon atom to which they are attached represent > C = O, > C = CH₂ or > C = NOCH₃.

Particularly important compounds of formula (I) which may be prepared according to the process of the present invention are those in which:

20 R¹ is isopropyi, R² and R³ together with the carbon atom to which the are attached represent >C = NOCH₃ and R⁴ is a hydrogen atom;

R' is isopropyl and R2, R3 and R4 are hydrogen atoms; and

R' is isopropyl, R2 is hydroxy and R3 and R4 are hydrogen atoms.

Compounds of formula (II) in which R¹ is a methyl, ethyl or isopropyl group, R² is a hydroxyl group and s R² is a hydrogen atom may are described in British Patent Specification 2187742A. The other compounds of formula (II) are described in European Patent Specification 0238258-A1.

The following Preparations and Examples illustrate the invention in which all temperatures are in °C and °L-" refers to Litre. In the following Examples hpic (high performance liquid chromatography) was carried out on a column of Spherisch SSODS-2 (150mm x 2.1mm) using acetonitrile/water (3:1) as eluant at a rate of 20 .5mm/min unless otherwise stated. The column effluent was interfaced with a Finnigan morphism belt to a Finnigan MAT4500 mass spectrometer under EL. conditions for mass spectrum characterisation. The compounds of the following Preparation are named by reference to the known parent 'Factor', Factor A. Factor A (described in UK Patent Specification 2168436) is a compound of formula (f) in which R¹ is an isopropyl group, R² is a Hydroxyl group and R² and R⁴ are hydrogen atoms.

## Intermediate 1

#### 40 5-Keto Factor A,

prepared as described in Example 2 of British Patent Specification 2187742A.

Intermediate 2

45

#### 5.23-Diketo Factor A,

prepared as described in Example 2 of EP-A1 238258.

Intermediate 3

### 5-Keto Factor A 23-ethyl ether,

prepared as described in Example 7 of EP-A1 238258.

#### Intermediate 4

#### 5-Keto Factor A 23-hemioxalate

5-Keto Factor A (330mg), finely divided calcium carbonate (200mg) and oxalyl chloride (0.07ml) were stirred together in dry dichloromethane (5ml) for 10 mln. 2N Hydrochloric acid (4ml) was then added followed 5 mln later by ethyl acetate. The organic phase was collected, dried and evaporated and the residue dissolved in diisopropyl ether. Dilution of this solution with petroleum ether (b.p. 40-60°) precipitated the title compound as a white flocculent solid (262mg), [a§\* +120° (c 0.6, CHG4).

# λEtOH

240nm (E<sup>'</sup> 425),  $_{'}$ max (CHBr<sub>3</sub>) 3440 (OH), 1800 (CO<sub>2</sub>H dimer), 1720 (ester), and 1676cm<sup>-1</sup> ( $\alpha$ , $\beta$ -unsaturated ketone),  $\delta$  (CDCl<sub>3</sub>) include 6.52 (narrow m, 1H), 5.07 (q,3Hz,1H), 3.78 (s,1H), 1.83 (s,3H), and 0.71 (d,7Hz,3H).

## Intermediate 5

10

### 5-Keto 23[E]-methoxyimino Factor A,

prepared as described in Example 16 of EP-A1 238258.

#### Example 1

25

(a) A spore suspension (0.1ml) of Streptomyces thermoarchaensis NCIB 12213 [prepared as described for Streptomyces thermoarchaensis NCIB 12015 in Example 2 of British Patent Specification 2187742A] was used to inoculate a flask containing the Medium A (2.5ml):

Medium A	gL-1
D-Glucose	2.5
Malt dextrin MD 30E (Roquette (UK)Ltd)	25.0
Arkasoy 50 (British Arkady Co.Ltd)	12.5
Molasses	1.5
K <sub>2</sub> HPO₄	0.125
Calcium carbonate	1.25
MOPS (3-(N-morpholino)propanesulphonic acid)	21.0
[Distilled water to 1 L, pH adjusted to 6.5 with 5N before autoclaving].	NaOH

This culture was incubated at 31 \* for 4 days on a rotary shaker operating at 250 rev/min, at which time intermediate 1 (1mg) in dimethylsulphoxide (0.05ml) was added. The culture was incubated for a further 24h at 31 \*, then centrifuged (approx. 10000 rpm/10 min), the supernatant removed and (1.8ml) of methanol added to the residue. The resulting suspension was allowed to stand for 1h at room temperature and then centrifuged. The supernatant was analysed by high exhibe showed by comparison with an authentic sample described in UK Patent Specification 2166436, the presence in the test sample of the compound of formula (1) where R¹ is an isopropyl group, R² is a hydroxyl group, R³ is a hydrogen atom and R¹ is a hydrogen atom (3.3% - supressed as % of the starting material added that was converted to the product).

(b) A repeat of the part (a) experiment with <u>Streptomyces avermittlis</u> ATCC 31780 and Intermediate 1 yielded an extract which on hpic analysis showed the presence of the compound of formula (I) in which R<sup>1</sup> is an isopropyl group, R<sup>2</sup> is a hydroxyl group, R<sup>3</sup> is a hydrogen atom and R<sup>4</sup> is a hydrogen atom [8.6% - % expressed as in part (a)].

(c) In a similar manner to part (a) Intermediate 1 was incubated with Streptomyces thermoarchaensis NCIB 12212, except that the culture was prepared from a spore suspension added to Medium A but which

after 3 days incubation as described had been used (0.1ml transferred) to inoculate a fresh portion of the same medium (2.5ml). Intermediate 1 (1mg in 0.05ml dimethylsulphoxide) was added to this later culture 2 days after and the culture was then incubated for a further two days before being harvested, extracted and analysed as described in part (a). Hplc analysis of a test sample showed the presence of the compound of formula (1) in which R¹ is an isopropyl group, R³ is a hydroxyl group, R³ is a hydrogen atom and R⁴ is a hydroxyl more fi.5 h²s - % expressed as in part (a)).

## Example 2

10

35

55

(a) A spore suspension (0.4ml) of <u>Streptomyces</u> thermoarchaensis NCIB 12213 [prepared as described for Example 1] was used to incoulate a <u>250ml</u> shake flask containing <u>Medium</u> A (25ml). This culture was incubated at 28° for 2 days on a rotary shaker operated at 250rev/mln. After <u>2</u> days a portion (5ml) of this culture was added to a 50ml shake flask and a 20mg/ml solution of Intermediate 2 in methanol (50µl) was added. The culture was incubated for a further 2-3 days under the same conditions, then treads with an equal volume of methanol. The resulting suspension was shaken for 1h and centifuged. The supernatant was analysed by hjc/mass spectrometry. A peak with a retention time 3 min. had a mass spectrum characteristic of the compound of formula (i) in which R¹ is isopropyl, R² and R³ together with the carbon atom to which they are attached represent >C = 0 and R⁴ is a hydrogen atom by comparison with an authentic sample described in UK Patent Specification 2176182.

(b) A repeat of the part (a) experiment with <u>Streptomyces thermoarchaensis</u> NCIB 12213 and Intermoidate 3 yielded an extract which on hplc/mass spectrometry analysis (using acetonitrile/water 9:1 as eluant) showed the presence of a compound of formula (i) in which R¹ is an isopropyl group, R² is an eluant) showed the presence atom and R⁴ is a hydrogen atom by comparison with an authentic sample described in UK Patent Specification 2176182.

(c) A repeat of the part (a) experiment with Streptomyces thermoarchaensis NCIB 12213 and Intermediate 4 yielded an extract which on hplc analysis showed the presence of a compound of formula (l) in which R an isopropyl group, R² is a group -OCOCo₂H, R³ is a hydrogen atom and R⁴ is a hydrogen atom by comparison with an authentic sample described in UK Patent Specification 2176182.

(d) A repeat of the part (a) experiment with Streptomyces thermozrchaensis NCIB 12213 and Intermediate 5 yielded an extract which an hpic analysis showed the presence of a compound of formula (l) in which R¹ is an isopropyl group, R² and R³ together with the carbon atom to which they are attached represent C = NOCH₃ and R¹ is a hydrogen atom by comparison with an authentic sample described in UK Patent Specification 2192630A.

E.I. mass spectroscopy of a sample isolated from analytical hplc yielded a molecular ion at 639 and gave characteristic fragments at 621, 527, 511, 496, 448, 434, 368, 264, 248 and 151.

#### Claims

10

20

30

25

1. A process for the production of compounds of formula (I)

[wherein R¹ is a methyl, ethyl or isopropyl group; R² is a hydrogen atom or a group  $OR^5$  (where  $OR^3$  is a hydroxyl group or a substituted hydroxyl group having up to 25 carbon atoms) and R³ is a hydrogen atom, or R² and R³ together with the carbon atom to which they are attached represent  $> C = > C = CH_2$  or  $> C = NOR^6$  (where R⁵ represents a hydrogen atom,  $C_{1-8}$  alkyl group or a  $C_{2-8}$  alkenyl group and the group  $> C = NOR^6$  is in the E configuration); and R⁵ is a hydrogen atom or a methyl group] which comprises incubating a compound of formula (II)

(wherein RI, R2 and R3 are as defined above) in a suitable medium in the presence of a microorganism or an enzyme derived therefrom or a preparation derived from a microorganism containing the enzyme of interest capable of effecting the conversion.

2. A process according to claim 1 in which the microorganism is a strain of Streptomyces thermoarchaensis, Streptomyces cyaneogriseus noncyanogenus, Streptomyces avermitilis or Streptomyces hygroscopicus subsp. aureolacimosus.

3. A process according to claim 1 in which the microorganism is Streptomyces thermoarchaensis NCIB 12212 and 12213 and mutants of these strains, Streptomyces avermitties ATCC 31272 or 31780 or Streptomyces hygroscopicus subsp. survolacimosus FERM P 1439.

- 4. A process according to claim 1 which comprises feeding the compound of formula (II) in a suitable souther into a fermentation medium comprising the microorganism in the presence of assimilable sources of carbon, nitrogen and mineral salts.
- 5. A process according to claim 1 which is carried out at a pH of 5.5 to 7.5 and a temperature of 25 to 5  $40^{\circ}$  C.
  - A process according to claim 1 in which R¹ in formula (I) and formula (II) is an isopropyl group.
  - 7. A process according to claim 1 in which, in formula (i) and formula (ii),  $R^2$  is a hydroxy or ethoxy group and  $R^3$  is a hydrogen atom; or  $R^2$  and  $R^3$  each represents a hydrogen atom; or  $R^2$  and  $R^3$  together with the carbon atom to which they are attached represent > C = 0, > C = C + 10, > C = R NOCHs.
- 8. A process according to claim 1 in which, in formula (I):
  - R¹ is isopropyl, R² and R³ together with the carbon atom to which they are attached represent > C = NOCH₃ and R⁴ is a hydrogen atom;
  - R1 is isopropyl, R2, R3 and R4 are hydrogen atoms; or

20

25

30

25

40

50

55

R' is isopropyl, R2 is hydroxy and R3 and R4 are hydrogen atoms.

## (12)

## EUROPEAN PATENT APPLICATION

- (2) Application number: 89302442.2
- (a) Int. Ci.5: C12P 17/18, //(C12P17/18, C12R1:465),A61K31:00

- 2 Date of filing: 13.03.89
- (39) Priority: 14.03.88 GB 8805978
- Date of publication of application:
  20.09.89 Bulletin 89/38
- Designated Contracting States:
   AT BE CH DE ES FR GB GR IT LI LU NL SE
- Date of deferred publication of the search report: 07.11.90 Bulletin 90/45
- Applicant: AMERICAN CYANAMID COMPANY
   One Cyanamid Plaza

- Wayne New Jersey 07470(US)
- (iii) Inventor: Rudd, Brian Arthur Michael 72 Exeter Road Rayners Lane Harrow Middlessex(GB) Inventor: Ramsay, Michael Vincent John 157 Kings Road South Harrow Middlesex(GB)
- Representative: Skalles, Humphrey John et al Frank B. Dehn & Co. Imperial House 15-19 Kingsway London WC2B 6UZ(GB)
- Process for the preparation of macrolide compounds.
- (f) S541 (LL-F28249) compounds of formula (I)

[wherein R¹ is a methyl, ethyl or isopropyl group; R² is a hydrogen atom or a group  $OR^5$  (where  $OR^5$  is a hydrogyl group or a substituted hydroxyl group) and  $R^3$  is a hydrogen atom, or  $R^3$  and  $R^3$  together with the carbon atom to which they are attached represent S=0, S=0, S=0, S=0. Then S=0 is the S=0 is a hydrogen atom, S=0 is a hydrogen atom or a methyl group and the group S=0. Then S=0 is the S=0 is the S=0 is a hydrogen atom or a methyl group) which comprises incubating a corresponding

5-keto compound in a suitable medium in the presence of a microorganism or an enzyme derived therefrom or a preparation derived from a microorganism containing the enzyme of interest capable of effecting the conversion.



## EUROPEAN SEARCH REPORT

Application Number

EP 89 30 2442

DOCUMENTS CONS	IDERED TO BE RELEVAN	T		
Category Citation of document with of relevant p	indication, where appropriate, passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)	
A EP-A-0 194 125 (M	ERCK AND CO., INC.)		C 12 P 17/18	
A EP-A-0 214 739 (M	ERCK AND CO., INC.)		(C 12 P 17/18 C 12 R 1:465)	
			A 61 K 31:00	
			TECHNICAL FIELDS	
			SEARCHED (Int. Cl.4)	
			C 12 P A 61 K	
	,,			
The present search report has	been drawn up for all claims			
Place of search	Date of completion of the search		Examiner	
THE HAGUE	21-08-1990	Novo	A Y SANJURJO M.A.	
CATEGORY OF CITED DOCUME	E - portion notant doc	umeot, but publi:	invention shed on, or	
X: particularly relevant if taken alone Y: particularly relevant if combined with an document of the same category	other D: document cited in	after the filing date D: document cited in the application D: document cited for other reasons		
A : technological background O : non-written disclosure P : intermediate document			, corresponding	